

### AGROMET ADVISORY BULLETIN

# GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI DEPARTMENT OF AGRONOMY, MPKV, RAHURI.



Ph. No. 02426 243239

243239 E-mail: rahuri.amfu@gmail.com

#### 58) Weather based Agromet Advisory committee meeting dated 17.09.2021

**District: Ahmednagar** 

Last Week Weather Summary							Weather	Weather Forecast				
(10.09.2021 to 16.09.2021)							Parameters	(18.09.2021 to 22.09.2021)				
10	11	12	13	14	15	16	Date	18	19	20	21	22
0	0	3.6	2.2	1.8	0	0	Rainfall (mm)	3	1	6	9	5
29.8	31	30.8	29.2	29.8	28.4	29.6	Max. Temp. (0C)	30	31	31	30	29
23.4	24.4	23.9	24.1	23.7	24.3	23.4	Min. Temp. (0C)	22	22	23	23	22
0	0	2	2	2	2	0	<b>Cloud Cover</b>	7	7	7	6	8
85	81	87	89	85	81	80	Max. RH (%)	87	87	86	87	86
58	65	76	64	67	59	57	Min. RH (%)	57	52	51	59	64
5	4	6.2	4.8	5.1	5.1	3.8	Wind Speed(km/hr)	10	11	12	13	15

**Agromet Advisory Based on Weather Forecast Prediction** 

Agromet Advisory Based on Weather Forecast Prediction							
Crop	Stage	Advisory					
Weather Summary		Sky will remain partly cloudy during next five days.					
		Very light rain is expected during next five days at isolated places.					
General A	Advisory	1) In case of manual field operations of harvesting/ picking, accomplish the					
		operation in 4-5 feet spaced strips assigning one strip to one person. This will					
		ensure adequate social distance to engaged labours. Agricultural machinery					
		should be sanitized frequently. Farmer should use face mask.					
		2) Farmers should take necessary precautionary measures while spraying					
		insecticide, fungicide etc.					
		3) Use "Meghdoot" mobile app for weather based crop advisory and prediction.					
CMC A	ly dia a my	Cotton: Spray the crop with Lambda-Cyhalothrin 5% EC 8 ml or Spinosand 45					
SMS Advisory		SC 3.5 ml per 10 liters of water for the control of Pink bollworm.					
Cotton	Opening	Spray the crop with Lambda-Cyhalothrin 5% EC 8 ml or Spinosand 45 SC 3.5					
	Boll	ml or Profenofos 50 EC 30 ml or Indoxacarb 15.8 EC 10 ml per 10 liters of					
		water for the control of Pink bollworm.					
Soybean	Pod	Spray the crop with Thiacloprid 29.7 SC 15 ml per 10 liters of water for the					
_	Formation	control of Girdle Beetle.					
Maize	Milking	Spray the crop with Thymethoxam 12.6% + Lambda Cyhalothrin 9.5% ZC at					
	_	0.5 ml per liter of water Or chlorantraniliprole 18.5% SC 4 ml per liter of water.					
		When spraying, care should be taken that the chemical does fall into the whorl					
		so that the effect of the chemical is seen properly.					
Pigeon pea	Branching	In pigeon pea if the infestation of stem fly is observed then spraying of 5%					
		NSKE.					
		As soon as incidence of sterility mosaic disease is noticed in pigeon pea crop,					
		uproot and destroy the diseases affected plants. The disease spreads through					
		mites. To control the mites, spraying of chemicals should be done.					
Groundnut	Harvesting	Harvest should be done of timely sown crop.					
Per millet	Harvesting	Harvest should be done of timely sown crop.					
Chickpea		Land preparation should be done for sowing of <i>rainfed</i> chickpea crop.					
		Collect and destroy the stubbles of kharif crop. Sowing of rainfed chickpea					
		crop should be done from 20th September onwards.					



### AGROMET ADVISORY BULLETIN

# GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI DEPARTMENT OF AGRONOMY, MPKV, RAHURI.



E-mail: rahuri.amfu@gmail.com

#### Ph. No. 02426 243239

Onion	Spray the crop with Lambda-Cyhalothrin 5% EC 6 ml + Tebuconazole 10 ml +
	stickers 10 ml per 10 liters of water for the control of thrips and blight. Spray
	should be done with simple hand pump.
Brinjal	The incidence of stem borers and fruit borers is likely to increase in Brinjal.
	Remove the infected plants and fruits and destroy it.
	Spray the crop with Cypermethrin 10% solution 10 ml Or Indoxacarb 14.5 EC
	10 ml Or Spinosad 45 EC 5 ml or Chlorantraniliprole 18.5 SC 3 ml per 10 liters
	of water for larval control. Spraying should be done with simple hand pump.
	Occasionally spray 5% neem extract.
Tomato	For control of sucking pest, spray Imidacloprid 5 ml or Thiomoxadam 4 gm per
	10 liters of water with a simple hand pump.
	Spray the crop with Cypermethrin 10 EC 10 ml + Trizophos 35% EC
	[compound pesticide] 20 ml + Tebuconazole 10 ml per 10 liters of water for
	control of fruit borer. Spray should be done with simple hand pump. Also
	occasionally spray 5% neem extract.

#### Source:

1) Weather Forecast : RMC, Mumbai

2) Last week weather summary : IMD Observatory, WMP, MPKV, Rahuri.

Place: MPKV, Rahuri Sd/-

Date : 17.09.2021 Nodal Officer, GKMS, AMFU Rahuri & Head,

Department of Agronomy, MPKV, Rahuri.