

AGROMET ADVISORY BULLETIN

GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI DEPARTMENT OF AGRONOMY, MPKV, RAHURI.



Ph. No. 02426 243239

E-mail: rahuri.amfu@gmail.com

81) Weather based Agromet Advisory committee meeting dated 06.12.2024

District: Dhule

| Last Week Weather Summary (29.11.2024 to 05.12.2024) | | | | | | Weather Parameters | Weather Forecast (07.12.2024 to 11.12.2024) | | | | | |
|--|----|----|----|----|----|-----------------------|---|----|----|-----|-----|-----|
| 29 | 30 | 01 | 02 | 03 | 04 | 05 | Date | 07 | 08 | 09 | 10 | 11 |
| | | | | | | | Rainfall (mm) | 0 | 0 | 0 | 0 | 0 |
| | | | | | | | Max. Temp. (0C) | 28 | 31 | 29 | 28 | 26 |
| | | | | | | | Min. Temp. (0C) | 15 | 14 | 15 | 16 | 15 |
| | | | | | | | Cloud Cover | 3 | 3 | 3 | 3 | 3 |
| | | | | | | | Max. RH (%) | 76 | 75 | 76 | 75 | 76 |
| | | | | | | | Min. RH (%) | 60 | 60 | 61 | 60 | 61 |
| | | | | | • | | Wind Speed(km/hr) | 1 | 3 | 4.5 | 4.1 | 6.5 |

Agromet Advisory Based on Weather Forecast Prediction

| | | Agromet Advisory Based on Weather Forecast Prediction |
|-------------------|---------|--|
| Crop | Stage | Advisory |
| Weather Summary | | According to the weather forecast from the RMC, Mumbai, sky will remain |
| | | cloudy during next five days. |
| General Advisory | | Use Meghdoot mobile app for weather-based crop advisory and prediction. |
| SMS Ad | visory | Sky will remain cloudy during next five days. |
| Chick pea Seeding | | |
| | | the monitoring of the gram pod borer install 5 pheromone trap/ ha above 1 m |
| | | from ground level. During the vegetative growth of crop take two preventive |
| | | spray at 15 days of interval of NSE 5% or Azadirachtin 0.30% 300 PPM @ 50 ml |
| | | per 10 liter of water. After the ETL level crosses by pod borer the spray |
| | | Emamectin benzoate 05 % SG @ 4.4 g or Emamectin benzoate 01.90 % EC 7.5 |
| | | ml or Indoxacarb 14.50 % SC @ 8-10 ml OR Indoxacarb 15.80 % EC @ 7 ml |
| | | OR Lambda-cyhalothrin 05 % EC @ 12.5 ml OR Spinetoram 11.70 % SC @ 1 |
| | | ml or Flubendiamide 8.33 % + Deltamethrin 5.56 % w/w SC @ 5 ml or |
| | | Novaluron 5.25 % + Indoxacarb 04.50 % SC @ 17 ml per 10m liter of water. |
| Wheat | CRI | Due to the cloudy weather there is a possibility of infestation of aphid and jassid, |
| | | spray the crop with 1 g Thiamethoxam 25WG in 10 liters of water as soon as |
| | | incidence of aphid and jassid noticed at ETL. |
| Safflower | Seeding | |
| | | and also to manage the weed flora, the intercultivation/ hoeing is to be carried out |
| | | at 25 to 30 DAS |
| Vegetables | | Due to cloudy weather, there is a possibility of increasing the prevalence of |
| crops | | sucking pest. For control of sucking insects, spray with Dimethoate 30 EC 15 ml |
| | | or Imidacloprid 17.8 SL 4 ml per 10 liters of water. |
| Tomato | | At initial incidence of Pinworm, install 16 yellow sticky traps per acre. In heavy |
| | | incidence, spray with Chlorantraniliprole 18.5 SC @ 3 ml or Flubendiamide 480 |
| | | SC @ 3 ml in 10 liters of water. |
| Brinjal | | Regularly monitor for shoot and fruit borer by using 5 pheromone traps per acre. |
| | | Infested shoots and fruits should be collected and buried into the soil. In heavy |
| | | incidence, spray with Chlorpyriphos 20 EC @ 20 ml or Emamectin benzoate 5 |
| ~ | | SG @ 4 g or Thiodicarb 50 WP @ 15 g in 10 liters of water. |
| Chili | | For management of leaf curl virus, remove infected plant part and spray |
| | | Acetamiprid 20 SP@ 3g per15 liters of water for control of whitefly vector. |
| Animal | | The rise in dust due to cold weather has resulted in a slight temperature increase, |
| Husbandry | 1 | which poses a greater risk for viral diseases. It's crucial to keep animals dry |
| | | during the occasional rains. With the sugarcane harvesting season underway and |



AGROMET ADVISORY BULLETIN

GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI DEPARTMENT OF AGRONOMY, MPKV, RAHURI.



E-mail: rahuri.amfu@gmail.com

Ph. No. 02426 243239

| | heightened animal transport, it is essential to prioritise vaccination against foot- | | | | | |
|--------|--|--|--|--|--|--|
| | and-mouth disease. There is a risk of pneumonia during this season. | | | | | |
| Mango | Spray Imidacloprid 17.8 SL @ 3 ml or Thiamethoxam 25 WG @ 4 g or | | | | | |
| | Buprofezin 25 SC @ 20 ml or Deltamethrin 2.8 EC @ 5 ml or Lambda- | | | | | |
| | cyhalothrin 5 EC@ 10 ml per 10 liters of water on entire mango tree for control | | | | | |
| | of mango hopper before flowering. | | | | | |
| Castor | For control of hairy caterpilar and spodoptera, spray Bacillus thuringiensis 0.5 | | | | | |
| | WP @ 20 g per 10 liters of water. In heavy incidence, spray Dimethoate 30 EC @ | | | | | |
| | 20 ml or Malathion 50 EC @ 20 ml 10 liters of water. | | | | | |

Source:

1) Weather Forecast : RMC, Mumbai

2) Last week weather summary :

Place: MPKV, Rahuri Sd/-

Date : 06.12.2024 Nodal Officer, GKMS, AMFU Rahuri & Head, Department of Agronomy, MPKV, Rahuri.