



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

Weather based Agromet Advisory committee meeting dated 26.02.2021

District: KOLHAPUR

Last Week Weather Summary (20.02.2021 to 26.02.2021)							Weather Parameters	Weather Forecast 27.02.2021 to 03.03.2021)				
20	21	22	23	24	25	26	Date	27	28	01	02	03
0.0	0.0	0.0	0.0	0.0	0.0	0.0	Rainfall (mm)	0	0	0	0	0
28.8	29.9	29.8	30.5	33.5	34.2	33.8	Max. Temp. (°C)	35	36	36	35	34
16.9	16.4	14.2	14.0	18.4	18.0	17.8	Min. Temp. (°C)	22	22	23	22	22
0	2	2	3	3	2	3	Cloud Cover	2	0	2	2	2
1	1	2	0	1	2							
76	88	88	82	96	89	85	Max. RH (%)	47	47	31	70	85
50	51	67	64	33	34	-	Min. RH (%)	23	14	11	14	19
4.6	4.6	4.8	10.2	6.2	6.1	-	Wind Speed(km/hr)	7	10	8	5	5
NE	E	N	NE	SE	W	W	Wind Direction	ESE	NNE	ENE	SE	ESE
E	SW	N	E	E	NE	-						
Rainfall previous week (mm)				Rainfall since 01.01.2021 (mm)			Rainy days since 01.01.2021					
0.0				24.8			2					

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary/ Alert		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is possibility of mainly clear weather from 27 th February to 03 rd March in Kolhapur district. Maximum temperature will remain nearby 34.0 to 35.0°C while minimum temperature will remain nearby 22 to 23°C. Wind speed will remain up to 05 to 10 km/hr.
General Advisory		<ul style="list-style-type: none"> ➤ Farmers should download "Meghdoot" android application on mobile for Agromet Advisory Services and "Damini" android application for forecast of lightening. ➤ Make arrangement of irrigation to the fruit orchards and rabbi crops as per the requirement.
SMS Advisory		Carry out harvesting and threshing of matured Wheat, Rabbi Sorghum and Chickpea crop. Dry the farm produce under sunlight and store at protective place.
Wheat	Maturity	➤ Carry out harvesting of early matured wheat varieties 2-3 days before full maturity of crop as it will avoid shattering of grains in field. At the time of harvesting moisture percentage in grain should be 15%.
Rabbi Sorghum	Maturity	➤ Carry out harvesting matured rabbi Sorghum crop. Dry the earheads under sunlight before threshing. Dry harvested produce/grains under sunlight and store at protected place.
Chick pea	Pod filling to maturity	➤ Chick pea crop matures within 110 to 120days. When pods get well dried carry out harvesting and then threshing of Chickpea crop. Dry the harvested chickpea grains under sunlight for 6 to 7 days and then store this dried produce at protected place. Keep the Neem leaves (5%) in stored grains so that it will prevent further insect damage to grains.
Sugarcane	Planting	➤ For ratoon management after harvesting of Sugarcane crop keep the sugarcane trash as it is, do not burn Sugarcane trash/leaves in field. Do not take ratoon if crop and field is infested with pest, disease and weed.



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

		<ul style="list-style-type: none">➤ Keep the ratoon of the Sugarcane crop which is harvested before 15th February. Otherwise there will be chances of Sugarcane stem borer.➤ Uproot and completely destroy grassy shoot of Sugarcane if found in Sugarcane ratoon crop.
	Preseasonal	<ul style="list-style-type: none">➤ During February month the rate of evaporation increases up to 5.49 mm per day. Sugarcane crop needs 3.27 liters of water per day thus for fulfilling that requirement Drip Irrigation Unit(4liters dripper) need to run for 49 minutes daily.
Onion	Growth	<ul style="list-style-type: none">➤ For control of Thrips in Onion take the spray of Dimethoate 30 % E.C. @ 15 ml or Lambda Cyhalothrin 5 % E.C. 6 ml mixed in 10 liters of water.
	Maturity	<ul style="list-style-type: none">➤ Stop irrigating to rangada Onion 3 weeks before ready to harvest.
Sweet potato	Planting	<ul style="list-style-type: none">➤ Carry out sowing of Summer Sweet Potato if there is availability of sufficient irrigation water source. Use Varsha, Konkan Ashwini, Samrat and Kalmegh varieties for sowing.
Brinjal	Sowing	<ul style="list-style-type: none">➤ Complete the planting of Brinjal in February-March. Select Manari gota, Phule Harit, Phule Arjun varieties for cultivation.
Cashew	Flowering	<ul style="list-style-type: none">➤ For control of tea mosquito use Lambda Cyhalothrin 5 % 6ml in 10 liters of water.
Animals	Growth	<ul style="list-style-type: none">➤ Carry out sowing of Hybrid Napier grass. Use Phule Jayvant and Phule Gunvant varieties.➤ During winter season allow Goats for grazing only after Sunrise when there is no dew on grass because during morning hours dew get formed on grasses when Goats feed on it they may suffer from respiratory diseases.

Source:

1) Weather Forecast

: Research Section, Mumbai

2) Last week weather summary

: IMD observatory (ZARS,KOLHAPUR)

Place : ZARS, KOLHAPUR

Date : 26.02.2021

Sd/-

Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur