



# AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA  
Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA,  
NANDURBAR



Ph. No. 02564 260544

E-mail: kvk\_ndb@yahoo.com

## 72) Nandurbar District : Agromet Advisory committee meeting dated 06<sup>th</sup> December, 2024

Weather Parameters	Weather Forecast (07.12.2024 to 11.12.2024)				
	07	08	09	10	11
Date	07	08	09	10	11
Rainfall (mm)	0	0	0	0	0
Max. Temp. (°C)	29	29	28	28	29
Min. Temp. (°C)	16	14	11	10	9
Cloud Cover (Octa)	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy	Partly Cloudy
Max. RH (%)	76	75	76	75	75
Min. RH (%)	61	62	61	61	60
Wind Speed(km/hr)	2	1	2	2	8

### Agromet Advisory Based on Weather Forecast

<b>Next Five Days Weather Forecast Summary</b>		According to weather forecast given by India Meteorological Department, Regional Meteorological Center, Mumbai, Possibility of sky may Partly Cloudy during next five days. Temperature may ranges maximum temperature 28 to 29 Degree Celsius & minimum temperature 9 to 16 Degree Celsius.
<b>Extended Range Weather Forecast (ERFS)</b>		Possibility of Maximum temperature may below Normal & Minimum temperature may Normal between 11 <sup>th</sup> to 17 <sup>th</sup> December 2024 over Madhya Maharashtra.
<b>Crop</b>	<b>Stage</b>	<b>Agromet Advisory</b>
<b>General Advisory</b>		Use Meghdoot mobile app weather forecast based crop specific agromet advisories
<b>SMS Advisory</b>		Carry out intercultural operation for soil moisture conservation and weed management.
<b>Chickpea</b>	<b>Sowing</b>	Keep chickpea crop field weed free for better growth of the crop. The first hoeing should be done after 20 days after sowing and the second after one month for soil moisture conservation.
<b>Rabi Sorghum</b>	<b>Seedling</b>	Apply top dressing of 40 kg N / ha 30 days after sowing to crop. The first 30 to 35 days of the crop should be kept weed free by weeding or hoeing.
<b>Cotton</b>	<b>Boll Development To Picking</b>	Picking of cotton better to carry out in the morning, as the moisture in the air does not cause the crop residues, dried leaves not to stick to the cotton. Good quality cotton should be picked separately, then yellow colored and rotted cotton & place separately. Each variety of cotton should be stored separately. After picking, the cotton should be dried in the sun for 3-4 days and stored in a clean & dry place.
<b>Pigeon Pea</b>	<b>Pod Development</b>	For control of Pod fly, Pod borer and plume moth, spray Emamectin benzoate (5% SG) 0.3 ml per liter of water or Indoxicarb (14.5 SC) 0.7 ml per liter of water.
<b>Papaya</b>	<b>Fruit Development</b>	During winter, Schedule irrigation at interval of 10 days as per requirement by double ring method, sprinkler or drip irrigation.
<b>Banana</b>	<b>Fruit Development</b>	Give irrigation to crop during night in winter. For protection from cold, cover the banana bunch with white plastic bag with 6% porosity.
<b>Mango</b>	<b>Development stage</b>	In case of outbreak of fungal disease due to cold, the infected branches should be cut 2 to 3 inches below the affected area and burnt. Apply Bordeaux paste or copper



# AGROMET ADVISORY BULLETIN

**DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA  
Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA,  
NANDURBAR**



**Ph. No. 02564 260544**

**E-mail: kvk\_ndb@yahoo.com**

		oxychloride paste on the cut area. Spray Carbendazim 1 gm or Copper Oxychloride 3 gm per liter of water on branches and stems.
<b>Animal Husbandry</b>	-	As minimum temperature is decreasing increase feed by including energy and protein rich minerals. Respiratory diseases are more likely to occur with the onset of cold so special care should be taken of young calves.

**Source:**

- 1) Weather Forecast : India Meteorological Department, Regional Meteorological Centre, Mumbai
- 2) Past Week Weather: District Agromet Unit, Krishi Vigyan Kendra, Nandurbar

**Place : Krishi Vigyan Kendra, Nandurbar**

**Date : 06.12.2024**

**Sd/-**

**Nodal Officer, GKMS-DAMU,  
KrishiVigyan Kendra, Nandurbar**