

Husbandry

AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA, NANDURBAR



E-mail: kvk_ndb@yahoo.com

Ph. No. 02564 260544

76) Nandurbar District: Agromet Advisory committee meeting dated 20th December, 2024

Weather	Weather Forecast (21.12.2024 to 25.12.2024)					
Parameters						
Date	21	22	23	24	25	
Rainfall (mm)	0	0	0	0	0	
Max. Temp. (°C)	29	29	28	29	30	
Min. Temp. (°C)	12	12	14	15	15	
Cloud Cover (Octa)	Clear	Clear	Clear	Clear	Clear	
Max. RH (%)	63	62	63	62	63	
Min. RH (%)	12	12	14	15	15	
Wind Speed(km/hr)	1	4	6	2	4	

Agromet Advisory Based on Weather Forecast						
Naxt Five Days Weather Forecast Summary		According to weather forecast given by India Meteorological Department, Regional Meteorological Center, Mumbai, Possibility of Weather may dry during next five days. Temperature may ranges maximum temperature 28 to 30 Degree Celsius & minimum temperature 12 to 15 Degree Celsius.				
Extended Range Weather Forecast (ERFS)		Possibility of Maximum temperature may Below normal & Minimum temperature may Normal between 25 th to 31 st December 2024 over Madhya Maharashtra.				
Crop	Stage	Agromet Advisory				
General Advisory		Use Meghdoot mobile app weather forecast based crop specific agromet advisories				
SMS Advisory		Carry out intercultural operation for soil moisture conservation and weed management.				
Cotton	Picking	Carry out picking of cotton and stored in a clean & dry place. Don't take ratoon crop of cotton which helps in reducing pink bollworm infection in next season.				
Wheat	Sowing To Seedling	Keep field weed free for better growth & development. Carry out intercultural operations for soil moisture conservation.				
Chickpea	Seedling	Keep field weed free for better crop growth. The first hoeing should be done after 20 days after sowing and the second after one month for soil moisture conservation.				
Rabi Sorghum	Seedling	In Rabi sorghum, for soil moisture conservation first hoeing should be done three weeks after sowing helps in weed control and addition of soil to the crop.				
	Pod	Harvest timely sown mature pigeonpea crop. For control of Pod fly, Pod borer and				
Pigeon Pea	Development To Maturity	plume moth, spray Emamectin benzoate (5% SG) 0.3 ml per liter of water or Indoxicarb (14.5 SC) 0.7 ml per liter of water.				
Papaya	Fruit	During winter, Schedule irrigation at interval of 10 days as per requirement by double				
I upuj u	Development	ring method, sprinkler or drip irrigation.				
Banana	Fruit	Give irrigation to crop during night in winter. For protection from cold, cover t				
	Development	banana bunch with white plastic bag with 6% porosity.				
Animal Husbandry	-	As temperature fall observed, farm livestock should be protected from exposure to cold particularly during night time to protect from pneumonia. Protect it from cold winds by				

hanging curtain from outside the shed.



AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA, NANDURBAR



P	o	11	ltı	·v
•	v	u		. .

-

Protect poultry birds from cold winds by hanging curtain from outside the poultry shed. Use of electric bulbs for heat is advisable. Increase feed to poultry birds by including energy and protein rich minerals.

Source:

1) Weather Forecast: India Meteorological Department, Regional Meteorological Centre, Mumbai

2) Past Week Weather: District Agromet Unit, Krishi Vigyan Kendra, Nandurbar

Place : Krishi Vigyan Kendra, Nandurbar Sd/-

Date : 20.12.2024 Nodal Officer, GKMS-DAMU,

KrishiVigyan Kendra, Nandurbar