

AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA, NANDURBAR



Ph. No. 02564 260544

260544 E-mail: kvk_ndb@yahoo.com

95) Weather based Agromet Advisory committee meeting dated 26th February, 2021

District: Nandurbar

Weather Parameters	Weather Forecast for Next Five Days (27 th February, 2021 To 03 rd March, 2021)				
Date	27	28	01	02	03
Rainfall (mm)	0	0	0	0	0
Maximum Temperature (°C)	35	35	35	35	35
Minimum Temperature (°C)	18	19	20	19	19
Cloud Cover (Octa)	Clear	Clear	Clear	Clear	Clear
Maximum Relative Humidity (%)	38	41	34	30	13
Minimum Relative Humidity (%)	16	18	13	12	08
Wind Speed (km/hr)	8	8	7	8	6

Agromet Advisory Based on Weather Forecast

Crop	Stage	Agromet Advisory				
Weather Forecast Summary		 According to weather forecast given by India Meteorological Department, Regional Meteorological Centre, Mumbai, During next five days Maximum and minimum temperature may ranges between 35 degree Celsius and 18 to 20 degree Celsius respectively. As per Extended Range Weather Forecast (ERFS) maximum and minimum temperatures above average between 4th to 10th March, 2021 over Madhya Maharashtra region. 				
General Advisory		➤ Keep crops weed free by intercultivation on time. Carry out plant protection measures to protect the crop from pest and diseases infestations on time.				
SMS Advisory		> During next five days weather may dry over the district.				
Gram	Maturity To Harvesting Stage	➤ If the rainfed gram is mature, it should be harvested and threshed. The gram should be dried in the sun for 6-7 days and stored in a dry and well ventilated place. Do not harvest when the crop is moist.				
Wheat	Development Stage To Maturity	Late sown wheat crop should be irrigated 80 to 85 days after sowing at grain filling stage. Harvest timely sown mature crop harvest and store at safe place.				
Rabi Sorghum	Maturity Stage	➤ Irrigate late sown sorghum when crop is at grain filling stage. Harvest timely sown mature crop harvest and store at safe place.				

AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA, NANDURBAR



E-mail: kvk_ndb@yahoo.com

Ph. No. 02564 260544

	Maize should be harvested when the husk cover is yellowish			
Rabi Maize Harvesting	white and firm. Harvest and dry it in the sun for 2-3 days.			
Groundnut Germination Stage	Carry out Gap filling after germination immediately. Apply irrigation 4-5 days after sowing of groundnut so that the remaining seeds germinate. Then irrigate 10 to 12 times at intervals of 8 to 10 days as per soil type.			
Onion Early Maturity Stage	➤ For Rabi onion, Irrigation should be stopped 10 to 15 days before harvest. Harvest the crop after 50% of the leaves have fallen.			
Papaya Development Stage	For the control of Aphids spray Dimethoate 10 ml OR Neemark 10 ml per 10 liters of water at 15 days interval.			
Watermelon Development stage	➤ Apply 50 kg Nitrogen fertilizer to watermelon 30, 45 and 60 days after planting.			
Banana Fruit Development Stage	> Two lines of wind breaks at 2 m around the banana orchard should be laid at a distance. Shaveri or Gajraj grass etc. should be used for wind breaks. The garden should be fenced on all four sides.			
Mango Blooming to Fruiting Stage	Due to the increase in temperature and decrease in humidity, pea to betel-shaped fruits in mango orchards are likely to fall on ground. To falling of this fruiting, depending on the availability of water, irrigate field at weekly intervals to the stage of betel nut size.			
Poultry Bird Flu	 Do's in backyard poultry rearing: Avoid contact with migratory birds in your poultry. Wash birds water and food containers daily. Keep the poultry shed clean. Vaccinate the poultry birds regularly. If any of poultry birds get sick or die, report it immediately to the nearest veterinary officer. Don'ts in backyard poultry rearing: Avoid contact with poultry droppings and nasal discharge. Avoid handling poultry birds as much as possible, wash hands with soap if handled. Do not leave poultry birds water and food outdoors. Make sure that other species of birds / animals do not come in the area. 			
➤ For Weather Forecast and Agromet Advisory use "MEGHDOOT" Mobile application.				

Source:

1) Weather Forecast : Regional Meteorological Centre, Mumbai

2) Last week weather : NOT AVAILABLE



AGROMET ADVISORY BULLETIN

DISTRICT AGROMET UNIT, GRAMIN KRISHI MAUSAM SEWA Dr. HEDGEWAR SEVA SAMITI, KRISHI VIGYAN KENDRA, NANDURBAR



Ph. No. 02564 260544 E-mail: kvk_ndb@yahoo.com

Place : Krishi Vigyan Kendra, Nandurbar Sd/-

Date : 26.02.2021 Nodal Officer, GKMS-DAMU,

Krishi Vigyan Kendra, Nandurbar