



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

Weather based Agromet Advisory committee meeting dated 28.07.2020

District: SANGLI

Last Week Weather Summary (22.07.2020 to 28.07.2020)							Weather Parameters	Weather Forecast (29.07.2020 to 02.08.2020)				
22	23	24	25	26	27	28	Date	29	30	31	01	02
1.9	2.0	0.2	0.0	0.0	0.5	-	Rainfall (mm)	12	19	24	20	26
31.0	-	29.0	29.0	29.2	28.8	-	Max. Temp. (0C)	32	33	33	33	33
23.6	22.7	23.8	22.9	22.4	21.9	-	Min. Temp. (0C)	22	22	22	22	23
8	8	8	8	7	3	-	Cloud Cover	8	8	8	8	8
92	96	96	95	93	95	-	Max. RH (%)	90	90	90	89	89
74	-	84	85	79	85	-	Min. RH (%)	61	60	61	63	64
0.0	1.9	1.9	0.0	3.7	1.9	-	Wind Speed(km/hr)	10	11	10	14	14
-	-	-	-	-	-	-	Wind Direction	W	W	W	W	W

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary/ Alert		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is possibility of light to moderate rainfall at scattered places from 29 th July to 02 nd August in Sangli district. Maximum and minimum temperatures will remain 32.0 to 33.0 and minimum up to 22.0 to 23.0 °C respectively. Wind speed will remain up to 10 to 14 km/hr.
General Advisory		<ul style="list-style-type: none"> ➤ Carry out hoeing / Weeding in kharif crop. Advice regarding protection and prevent infection from Coronavirus (COVID 19) disease. ➤ While going to purchase agricultural inputs in market follow social distancing, always wear mask. Sanitize hands with sanitizer or wash hands frequently.
SMS Advisory		➤ Heavy rainfall is likely to occur from 28 th to 30 th July in ghat region of Sangli district and thunderstorm accompanied with lightning and heavy rainfall likely to occur on 1 st August at isolated places in district Sangli.
Maize	Vegetative growth	<ul style="list-style-type: none"> ➤ Carry out hoeing in early sown crop. ➤ If infestation of American Fall Army Worm observes more than 10 % in Kagal, Shirol, Hatkanangale and Karvir tahsil then carry out spraying of one of the insecticide mentioned below. Chlorantraniliprole 18.5% SC insecticide @0.4 ml per litre of water. Or Spinetoram 11.7% SC @0.5 ml per litre of water. Or Emamectin Benzoate 5% SG @0.4 gram per litre of water. ➤ If there is shortage of soil moisture then give protected irrigation.
Kharif Jowar	Vegetative growth	<ul style="list-style-type: none"> ➤ Carry out hoeing in early sown crop. ➤ If there is shortage of soil moisture in tahsils from eastern part of district then give protected irrigation.
Green gram and Black gram	Vegetative growth	<ul style="list-style-type: none"> ➤ Carry out 1st hoeing 20 to 25 days after sowing of crop. ➤ If there is shortage of soil moisture in tahsils from eastern part of district then give protected irrigation.
Groundnut	Vegetative growth	<ul style="list-style-type: none"> ➤ Frequent rains and high humid condition is favorable for occurrence of tikka and Rust disease to control these diseases sprays the crop with Mancozeb 25 gram + 25 gram Bavistin by mixing in 10 liters of water. ➤ For control of sucking pest spray the crop with the solution prepared by mixing Methyl Demeton 25 E.C. 10 ml in 10 liters of water.
Rajama	Vegetative growth	<ul style="list-style-type: none"> ➤ Carry out hoeing in early sown crop. ➤ Apply second dose of nitrogen @30kg (70 kg Urea) per hector at the time of 20 days after sowing.
Soybean	Vegetative growth	➤ High humidity and high wind speed favourable for occurrence of rust in soybean. Brown spot observes on leaves and leaves get brown coloured due to infection, grain yield and per hector



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

		production of Soybean get decreases. If disease occurs take the spray of Propiconazole @ 1ml in 1 liter of water.
Sugarcane	Adsali Planting	<ul style="list-style-type: none">➤ Carry out Sugarcane planting in Adasali seasons up to 15th August for better yield and sugar recovery.➤ Select the varieties Co.86032 (Nira) and Co.M.0265(Phule 265) for Adsali Sugarcane cultivation.➤ Planting of sugarcane with the 30 to 45 old plants grown from single bud in nursery is very beneficial compared to planting method with sets as it saves irrigation, weeding, fertilizer and supervision required during 30 to 45 days period.➤ For planting of Sugarcane with sets select 9 to 10 month old healthy, genetically pure sugarcane sets.➤ Mix Dimethoate 30 % EC 265 ml in 100 liters of water and deep Sugarcane sets into the prepared solution for 10 minutes which protects the Sugarcane form fungal diseases and mealy bugs.➤ Prepare the solution by mixing Acetobactor and Phosphorus solubilizing Bacterial culture 10 kg and 1.25 kg respectively in 100 liters of water. Deep the Sugarcane sets into the prepared solution for 10 minutes and then carry out planting which saves 50 % Nitrogenous and 25 % Phosphorus fertilizers dose.
Onion	Seedling	<ul style="list-style-type: none">➤ For planting of Rangada Onion carry out sowing of seeds on raised beds.(If there is availability of irrigation facility)
Animals	Growth	<ul style="list-style-type: none">➤ As there is possibility of rainfall keep the animals at safe places.➤ During rainy season do not give excess of green fodder to animals.➤ In rainy season wounds gets formed near hoofs of sheeps to prevent this prepare the solution of Copper Sulphate or Formaldehyde in tank up to the height that hoofs gets dip into the solution and allow the sheeps to go from that solution.➤ To prevent nuisance from ectoparasites spray Deltamethrin (Ex. Butox) by taking advice from Veterinary doctor in goat shade and on the goats.

Source:

1) Weather Forecast

: Research Section, Mumbai

2) Last week weather summary

: IMD observatory (ZARS,KOLHAPUR)

Place : ZARS, KOLHAPUR

Date :28.07.2020

Sd/-

Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur