



AGROMET ADVISORY BULLETIN

GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR

ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

Weather based Agromet Advisory committee meeting dated 13.10.2026

District: SANGLI

Last Week Weather Summary (07.10.2026 to 13.10.2026)							Weather Parameters	Weather Forecast (14.10.2026 to 18.10.2026)				
07	08	09	10	11	12	13	Date	14	15	16	17	18
-	-	-	-	-	-	-	Rainfall (mm)	0	0	3	6	10
-	-	-	-	-	-	-	Max. Temp. (°C)	35	35	35	33	32
-	-	-	-	-	-	-	Min. Temp. (°C)	23	23	25	25	23
-	-	-	-	-	-	-	Cloud Cover	2	2	4	5	7
-	-	-	-	-	-	-	Max. RH (%)	71	74	78	90	95
-	-	-	-	-	-	-	Min. RH (%)	53	53	57	63	69
-	-	-	-	-	-	-	Wind Speed(km/hr)	10	11	10	6	4
-	-	-	-	-	-	-	Wind Direction	E	ENE	ENE	NE	ESE

Agromet Advisory Based on Weather Forecast

Crop	Stage	Advisory
Weather Summary		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated there is forecast of mainly clear sky on 14 th and 15 th October and generally cloudy sky on 16 th , 17 th and 18 th October in Sangli district. There is forecast of light rainfall at isolated places on 16th, 17th and 18th October in Sangli district. During next 5 days and Tmax may remain between 32.0 to 35.0°C, minimum temperature may remain nearby 23.0 to 25.0. Morning relative humidity will remain nearby 71 to 95% and Afternoon relative humidity will remain nearby 53 to 69%. Wind speed will remain between 04 to 11 km per hour.
ERFS		According to the Extended Range Forecast (ERF), over Madhya Maharashtra during 16 th to 22 nd October rainfall is likely to remain normal and maximum temperatures likely to remain above normal and minimum temperatures are also likely to remain above normal.
General Advisory		➤ Use drip or sprinkler irrigation for irrigating Kharif crop as well as in fruit and vegetable crop. ➤ Spray insecticides and fungicides during a rain-free period and when the wind is calm. ➤ If possible Carry out mulching in fruits and vegetable crop ➤ As rainfall is likely from 16 to 18 October, harvest mature rice, maize, soybean and groundnut crops. Wherever possible, complete threshing and promptly store the produce in a dry and safe place.
SMS Advisory		➤ For the rabi season, assess water availability and select suitable crops and varieties accordingly.
Sugarcane	Vegetative growth	➤ Do not begin preparations for Preseasonal sugarcane planting unless sufficient irrigation water is available. ➤ Increased humidity and higher maximum temperatures have led to the incidence of woolly aphid at some locations. As soon as the infestation is noticed, spray Imidacloprid 17.8% SL @ 3 ml/10 lit of water.
Rabbi Sorghum	Sowing	➤ By considering water availability carry out sowing of rabbi sorghum at vafasa condition in soil at 5 cm depth, Spacing :45x15cm, ➤ Seed treatment: Apply 4 gram Sulphur (300 Mesh) and 25 gram Azatobactor and P.S.B. Culture to per kg of seed.
Maize	Grain filling/ Preparatory	➤ Provide protection irrigation to Maize crop as per the availability ➤ Carry out harvesting of matured crop and keep the harvested produce at safe place as there is forecast of rainfall from 15th to 17th October.



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

	tillage	➤ Start preparatory tillage for <i>Rabi</i> season Maize crop only if there is availability of irrigation
Soybean	Pod filling to maturity	➤ Harvesting of Soybean should carry out after pods gets yellowish red colour and after completion of 100 to 110 days from sowing as per the maturity of crop variety. If harvesting delays pods breaks and seeds gets shattered in field. Harvesting should carry out nearer to ground with help of sharp sickle. Keep the harvested produce at safe place ➤ For late sown and long duration varieties which are at the pod-filling stage, spray 00:00:50 water-soluble fertilizer @ 10 g per litre of water.
Groundnut	Maturity	➤ Groundnut leaves turns yellow, shell of pods become tough and inner side of shell gets blackish colour after attaining maturity. Carry out harvesting of matured crop. Harvested pods should be dried under sunlight as soon as sunlight gets available then store dried pods at dry and safe place. Do not store pods in wet and humid place.
Chick Pea	Presowing	➤ Collect the inputs required for sowing of <i>rabi</i> chickpea and Carry out sowing of <i>rabi</i> chickpea upto 10th October at sufficient moisture condition (<i>vafasa</i>) in soil. ➤ Use Vijay, Digvijay, Phule Vikram and Phule Vishvaraj varieties for sowing. ➤ Seed treatment: Treat the seeds <i>Trychodarma Harjiyanam</i> 1.0% W.P. 6 gram per kg of seeds for prevention from Neck rot and root rot then treat the seeds with 250 gram Rayzobium and P.S.B. per 10 kg of Seeds by mixing in cold Jaggary solution. Dry this treated seed under shade almost for 1 hour and then carry out sowing. ➤ For control of Gram Pod borer through Integrated Pest Management carry out sowing of 200 gram of Sorghum by mixing in Chick Pea.
Turmeric	Rhizome development	➤ During the past week, cloudy and humid weather has favoured the incidence of leaf blight in some areas. To control the disease, spray Azoxystrobin 18.2% + Difenoconazole 11.4% SC @ 10 ml per 10 litres of water.
Ginger	Rhizome development	➤ Rhizome Fly (Caterpillar/Maggot) Control: ○ Perform timely earthing-up (hilling up soil around the base). ○ Use 6 wide earthen or plastic pans per hectare. In each pan, add 200 g of crushed castor seeds and 1.5 liters of water. After 8 to 10 days, as the mixture begins to emit an odor, the adult rhizome flies are attracted to the trap and drown/die. Since this control measure is highly effective, low-cost, and easy to implement, it plays a vital role in organic ginger cultivation. It is especially effective because adult flies are eliminated before their larvae can damage the rhizomes.
Animals	Growth	➤ Provide vaccination to the healthy Cattles from authorized for prevention from Foot and Mouth disease ➤ If sufficient soil moisture is available, sow <i>rabi</i> sorghum for fodder to ensure an adequate fodder supply for livestock. If irrigation water is available, cultivate hybrid Napier grass. ➤ If fodder maize is available, use it to prepare silage.
To download “ Meghdoot ” app scan QR code		To download “ Damini ” app scan QR code
		

Source:

1) Weather Forecast: Research Section, Mumbai

Place : ZARS, KOLHAPUR

Date : 13.10.2026

Sd/-

Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur