AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR

Ph. No. 0231 2692416 **E-mail: adrkolhapur@rediffmail.com**

Weather based Agromet Advisory committee meeting dated 09.11.2026

District: SATARA

Last Week Weather Summary (03.10.2026 to 09.11.2026)							Weather Parameters	Weather Forecast (10.11.2026 to 14.11.2026)				
03	04	05	06	07	08	09	Date	10	11	12	13	14
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Max. Temp. (^{0}C)	34	33	33	32	32
-	-	-	-	-	-	-	Min. Temp. (^{0}C)	15	15	14	13	13
-	-	-	-	-	-	-	Cloud Cover	0	0	1	1	2
-	-	-	-	-	-	-	Max. RH (%)	68	67	67	69	73
-	-	-	-	-	-	-	Min. RH (%)	40	39	39	40	44
-	-	-	-	-	-	-	Wind Speed(km/hr)	7	5	5	5	6
-	-	-	-	-	-	-	Wind Direction	E	E	ENE	ENE	ENE

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary/ Alert		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is forecast of mainly clear sky on 10th, 11th, 12th, 13th and 14th November in **Satara district. During next 5 days Tmax may remain 32.0 to 34.0 °C and minimum temperature may decrease by 2 to 3 °C and remain between 13.0 to 15.0°C.** Morning relative humidity will remain nearby 67 to 73% and Afternoon relative humidity will remain nearby 39 to 44%. Wind speed will remain between 05 to 07 km per hour.
ERFS		According to the Extended Range Forecast (ERF), over Madhya Maharashtra during 11th to 17th November rainfall is likely to remain below normal and maximum temperatures likely to remain normal and minimum temperatures are also likely to remain normal.
General Advisory		➢ Use drip and sprinkler irrigation for irrigating fruit and vegetable crop ➢ Use Ridges and Furrow, Broad Bed Furrows for rabbi crop. ➢ After harvesting of Sugarcane crop collect the stubbles ➢ Minimum temperature is expected to decrease, so farmers should take care of themselves and their livestock.
SMS Advisory		➢ If possible Carry out mulching in fruits and vegetable crop by using sugarcane trash
Rice	**Maturity**	➢ Harvest mature short-duration and mid-late rice varieties. Dry the harvested crop for 3–4 days, then thresh it and store the threshed grain in a safe, dry place.
Maize	**Maturity/ Sowing**	➢ Carry out harvesting of matured crop and keep the harvested produce at safe place ➢ Harvest the matured crops and ensure that the harvested and threshed produce is stored safely. ➢ Farmers should undertake maize sowing for the Rabi season up to 15th November only if assured irrigation facilities are available. ➢ Select improved early-maturing maize varieties for sowing. Maintain a spacing of 75 × 20 cm between rows and plants. ➢ At the time of sowing, apply 35 kg urea, 150 kg Single Super Phosphate (SSP), and 28 kg Muriate of Potash (MOP) per acre. ➢ Before sowing, treat maize seeds with Chlorantraniliprole 50% FS @ 5 ml per kg of seed. ➢ Immediately after sowing or on the following day, apply the recommended pre-emergence herbicide Atrazine @ 500 gram mixed in 200 litres of water per acre as a uniform spray on the soil surface. ➢ Install 5 pheromone traps per acre within one week after crop emergence for monitoring fall armyworm infestation.
Wheat	**Sowing**	➢ Only when there is availability of assured irrigation then and then only farmers can carry out sowing of Wheat crop up to 10th November. ➢ Make availability of Phule Samadhan, Phule Anupam, Phule Satwik, M.A.C.S. 6222 and NIAW 1149 varieties for sowing.

AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR

Ph. No. 0231 2692416 **E-mail: adrkolhapur@rediffmail.com**

		➢ Seed rate: Wheat under protective irrigation require: 75 to 100 kg seed per hector. ➢ Seed Treatment: Treat the seed with Thiram 75 % WS @ 3gram per kg of seed also for control of Aphids, Thrips and Stem fly ➢ Treat the seeds with Thiamethoaxam 30% F.S.@ 7.5ml per 10 kg of seed. Carry out seed treatment with fungicide, insecticide and allow drying the seed and then treat the seed with Azatobactor@25gram per 1 kg of seed and PSB @ 25gram per 1 kg of seed. This will helpful to increase yield by 10 to 15%.
Chick Pea	**Sowing**	➢ Only if there is availability of sufficient irrigation then farmers can carry out sowing of irrigated chickpea up to 10th November ➢ Use Vijay, Vishal, Digvijay, Phule Vikram, Phule Vikrant and Phule Vishvaraj varieties for sowing. ➢ Seed treatment: Treat the seeds Trychodarma Harjiyanam 1.0% W.P. 6 gram per kg of sees for prevention from Neck rot and root rot then trat the seeds with 250 gram Rayzobium and P.S.B. per 10 kg of Seeds by mixing in cold Jaggary solution. Dry this treated seed under shade almost for 1 hour and then carry out sowing. ➢ For control of Gram Pod borer through Integrated Pest Management carry out sowing of 200 gram of Sorghum by mixing in Chick Pea.
Sugarcane	**Vegetative growth**	➢ Plant pre-seasonal sugarcane only if an adequate supply of irrigation water is assured. ➢ Considering water availability, apply the second dose of nitrogen to adsali sugarcane 6–8 weeks after planting, at the rate of 139 kg urea per acre, equivalent to approximately 3.09 bags of 45 kg each. ➢ Use drip irrigation for sugarcane and apply water-soluble fertilizers through the drip system. ➢ To conserve soil moisture, remove the lower dry leaves of sugarcane and use them as mulch. ➢ Irrigate sugarcane through alternate furrows. ➢ Do not burn sugarcane trash. Use it as mulch wherever possible, and apply a microbial decomposer culture to accelerate decomposition. This will add organic matter to the soil.
Strawberry	**Vegetative growth**	➢ To obtain higher yields in strawberry, spray Gibberellic Acid (GA_3) @ 25 ppm at 30 days after planting. ➢ To obtain higher yields in strawberry, apply 35:40:18 kg of Nitrogen, Phosphorus and Potassium (N:P:K) per hectare, respectively, during the second month after planting, divided into three equal splits.
Mango	**New flush**	➢ Due to partially cloudy and humid weather there is attack of Leaf hoppers in some parts thus to control spray the Mango crop with Imidacloprid 17.8 S.L. 3ml or Lambda Cyhalothrin 5% E.C. 10 ml per 10 liters of water.
Animals	**Growth**	➢ To protect livestock against foot-and-mouth disease (FMD), vaccinate healthy animals in consultation with a qualified veterinarian. ➢ Provide each animal with 50 g of mineral mixture per day. ➢ During cold weather, farmers should keep their animals in warm, dry shelters and protect them from cold winds. Provide sufficient feed, clean drinking water, and warm coverings when needed. Special care should be taken of young and sick animals.
Poultry	**Growth**	➢ During cold weather, poultry farmers should keep birds in warm, dry, and well-ventilated sheds. Protect birds from cold winds, provide sufficient feed and clean drinking water, and maintain proper heating, especially for young chicks. Keep the litter dry to prevent diseases.

To download **"Meghdoot"** app scan QR code	To download **"Damini"** app scan QR code

Source:
1) Weather Forecast : Research Section, Mumbai

AGROMET ADVISORY BULLETIN

GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR

ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR

Ph. No. 0231 2692416 **E-mail: adrkolhapur@rediffmail.com**

Place : ZARS, KOLHAPUR
Date : 09.11.2026

Sd/-
Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur