



# AGROMET ADVISORY BULLETIN

GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI  
DEPARTMENT OF AGRONOMY, MPKV, RAHURI.



Ph. No. 02426 243239

E-mail: rahuri.amfu@gmail.com

## 96) Weather based Agromet Advisory committee meeting dated 30.01.2026

**District: Dhule**

Last Week Weather Summary (23.01.2026 to 29.01.2026)							Weather Parameters	Weather Forecast (31.01.2026 to 04.02.2026)				
23	24	25	26	27	28	29	Date	31	01	02	03	04
							Rainfall (mm)	1	0	0	4	1
							Max. Temp. (0C)	29	30	31	31	30
							Min. Temp. (0C)	17	19	19	18	19
							Cloud Cover	5	5	2	5	2
							Max. RH (%)	60	57	51	52	54
							Min. RH (%)	27	25	22	22	22
							Wind Speed (km/hr)	7.2	3.6	6.1	5	7.6

### **Agromet Advisory Based on Weather Forecast Prediction**

Crop	Stage	Advisory
<b>Weather Summary</b>		According to the weather forecast from the RMC, Mumbai, <b>sky will remain partly cloudy during next five days. There is also a possibility of very light rain during January 31<sup>st</sup>.</b>
<b>General Advisory</b>		1) Use Meghdoot mobile app for weather based crop advisory and prediction. 2) Farmers should take necessary precautionary measures while spraying insecticide, fungicide etc. and use Kisan Kavach Body Suite.
<b>SMS Advisory</b>		<b>Wheat:</b> Spray the crop with Mancozeb 75% WP 30g per 10 liters of water or Tebuconazole 50% + Typhloxystrobin 25% WP 10g per 10 liters of water should be sprayed as soon as the infestation of rust disease appears.
<b>Rabi Sorghum</b>	<b>Harvesting</b>	Protect matured sorghum crop from Birds. At the maturity stage the tip of sorghum grain gets black spot. Carry out harvesting of matures crop. Keep the earheads under sunlight for 8 to 10 days for proper drying. Mature crops should be harvested and harvested crops should be threshed and stored in a safe place.
<b>Wheat</b>	<b>Milking</b>	Give irrigation to wheat crop 80 to 85 days after sowing at grain filling stage when wind is calm which helps in to prevent crop from lodging. Rust disease is likely to increase in wheat crop, Mancozeb 75% WP 30g per 10 liters of water or Tebuconazole 50% + Typhloxystrobin 25% WP 10g per 10 liters of water should be sprayed as soon as the infestation appears.
<b>Chick pea</b>	<b>Pod formation</b>	To monitor the incidence of pod borer, install 5 pheromone traps per hectare. Large numbers of moths get trapped in these, which helps suppress reproduction. Additionally, for control, spray 5% neem seed extract. Chemical Control: Apply either Emamectin Benzoate 5% SG at 4 grams per 10 liters of water or Quinalphos at 20 ml per 10 liters of water-one of these chemical insecticides.
<b>Maize</b>	<b>Tasseling</b>	<u>American fall army worm control measures:</u> Mechanical method: Eggs, larvae fall armyworms should be removed and destroyed by mixing them in kerosene. Collect and destroy large numbers of male moths using pheromone traps. <u>Chemical Control:</u> Spray the crop with Chlorantraniliprole 18.5% SC at 40 GAI/ha or Emamectin Benzoate 5% SG at 8 g per liter. Or



**AGROMET ADVISORY BULLETIN**  
**GRAMIN KRISHI MAUSAM SEWA, AMFU, RAHURI**  
**DEPARTMENT OF AGRONOMY, MPKV, RAHURI.**



Ph. No. 02426 243239

E-mail: rahuri.amfu@gmail.com

		<p>Spray the crop with Spinetoram 11.7 SC insecticide at 0.5 ml per liter of water. Or Spray the crop with Emamectin 5% + Benzoate Lufenuron 40% WG at 0.2 g or Novaluron 5.25% + Emamectin Benzoate 0.9% SC at 3 ml per liter of water.</p>
<b>Summer Groundnut</b>	<b>Pre sowing preparation</b>	<p>Pre-sowing irrigation should be done to ensure that sowing can be done in the third week of January. Procured Groundnut seed of improved varieties like Phule Unnati, T.A.G-24, T.G-26, T.P.J-41, JL-501 etc. Summer groundnut should be sown between 15<sup>th</sup> January and 15<sup>th</sup> February 2023 but to avoid sowing till temperature raise up to <b>or</b> above 18 degree Celsius. Prepare land for summer groundnut and apply FYM @ 5 ton/ha.</p>
<b>Sugarcane</b>	<b><i>Suru Sugarcane Cultivation</i></b>	<p><u>Suru Sugarcane Cultivation:</u> Select varieties like Co 86032 (Nira), Co M0265 (Phule 265), Phule 10001, Phule 11082, Phule Sugarcane 15012, Phule 13007, Co 94012 (Phule Savitri), Co 92005 for cultivation. <u>Sets Treatment:</u> Before planting, soak the Sets in a solution of 10 grams of Carbendazim mixed with 10 liters of water for 10-15 minutes. Then, mix 1 kg of <i>Azotobacter</i> and 125 grams of phosphate-solubilizing bacteria in 100 liters of water, soak the cuttings in it for 30 minutes, and plant them immediately.</p>
<b>Onion</b>	<b><i>Late Kharif (Rangada)</i></b>	<p>Harvest onions when 50% of the foliage (tops) has fallen over. If continuous leaf growth persists, roll the drum and harvest 15-20 days later. After harvesting, dry the onions in the field along with their tops for 4-5 days in rows. Trim the tops leaving 3-4 cm stubs. Grade the onions and send them for sale.</p>
<b>Onion</b>	<b><i>Rabi/Summer</i></b>	<p>For <i>Rabi/summer</i> onion planting, use flat beds. Prepare beds 2 meters wide and 3-5 meters long according to the land slope before forming ridges, apply well-decomposed FYM. Transplant seedlings at 15 x 10 cm spacing.</p>

**Source:**

1) Weather Forecast : RMC, Mumbai  
2) Last week weather summary :

**Place : MPKV, Rahuri**

**Sd/-**

**Date : 30.01.2026**

**Nodal Officer, GKMS, AMFU Rahuri &  
Head, Department of Agronomy, MPKV, Rahuri.**