



**AGROMET ADVISORY BULLETIN**  
**GRAMIN KRISHI MAUSAM SEWA, AMFU, PUNE**  
**Department of Agricultural Meteorology**  
**College of Agriculture, Pune 411 005**



Ph No. 020-29516264

E-mail: [amfupune@gmail.com](mailto:amfupune@gmail.com)

**Weather based Agromet Advisory committee meeting dated 05.12.2025**  
**District Jalgaon**

Weather Parameters	Weather Forecast (05.12.2025 to 09.12.2025)				
	Date	6	7	8	9
Rainfall (mm)	0	0	0	0	0
Max. Temp. (0C)	29	30	29	29	29
Min. Temp. (0C)	13	12	13	12	10
Cloud Cover	0	2	1	0	0
Max. RH (%)	66	59	55	51	56
Min. RH (%)	39	35	29	29	31
Wind Speed(km/hr)	9	8	7	7	7
Wind direction (deg)	86	80	96	100	84

**Agromet Advisory Based on Weather Forecast Prediction**

Crop	Crop Stage	Advisory
<b>Weather Summary/ Alert</b>		According to forecast given by Regional Meteorological Centre, Mumbai, India Meteorological Department, the weather may remain dry and cold during dt. 05 <sup>th</sup> to 09 <sup>th</sup> December, 2025 in the district.
<b>Extended Range Forecast (ERFS)</b>		As per ERFS products during 10 <sup>th</sup> to 16 <sup>th</sup> December, 2025 over Madhya Maharashtra (Dhule, Nandurbar, Jalgaon, Nashik, Ahmednagar, Pune, Satara, Sangli, Solapur, Kolhapur) division <ul style="list-style-type: none"> <li>➤ No Rainfall.</li> <li>➤ Maximum temperature may remain moderately below normal.</li> <li>➤ Minimum temperature may remain below normal.</li> </ul>
<b>General Advisory</b>		<ul style="list-style-type: none"> <li>• Apply irrigation to orchard/ crop in night hours as the minimum temperature is likely to remain low.</li> <li>• Do not apply fertilizer to crop during cold weather condition as plants can't uptake it efficiently due to poor root activity.</li> <li>• Farmers should observe the pest and disease incidence in the field and if the infestation is above economic threshold level (ETL), then proper pest and disease management should be done.</li> <li>• Farmers should spray insecticides and herbicides separately.</li> <li>• Before spraying any pesticide, fungicide or herbicide on any crop, farmers should make sure whether the product content is as per label claim.</li> <li>• While cultivating vegetable crops like brinjal, okra, guar etc., make use of yellow sticky traps, host trap crops, biological fungicides (Metarhizium, Trichogramma, Verticillium, etc.) should be used after planting according to the stage of the crop for the biological control of sap-sucking insects. Chemical insecticides and fungicides should be used if necessary.</li> <li>• Farmers should download and make use of 'Meghdoot' mobile app for weather based crop advisory and weather forecast.</li> <li>• Also, farmers should download and make use of 'Damini' mobile app for</li> </ul>

		<p>lightning and thunderstorm warning and forecasting.</p> <ul style="list-style-type: none"> <li>• Farmers should refer and use 'KRISHIDARSHINI' published by Mahatama Phule Agricultural University for all agriculture related information.</li> <li>• Micronutrient powder Phule Grade-I should be applied to the soil at the time of sowing. Phule Grade II is a liquid micronutrient mixture which is specially recommended for foliar application at 1% concentration. For this, the first spray should be done about 30-45 days after sowing. The second spray should be repeated 15-20 days after the first spray. The combination of soil application of Grade I and foliar spray of Grade II ensures immediate and sustained availability of micronutrients during the growth period of the crop, which leads to good growth, yield and quality of the crop. This should be used at sowing of <i>rabi</i> crops like sorghum, gram, wheat etc.</li> </ul>
<b>Wheat</b>	Vegetative Growth Stage	As per requirement and soil condition/type apply irrigation to the wheat crop 18 to 21 days after sowing at the time of crown root initiation.
<b>Cotton</b>	Picking & storage stage	When 30 to 35 % of bolls are opened, first picking of cotton should be carried out and there after interval of 15 to 20 days 2 to 3 pickings can be done. The harvested produce should be dried in sunlight for 3 to 4 days and kept at dry and clean places.
<b>Maize</b>	Grain filling stage	In standing crop, spraying of neem extract 5% or emamectin benzoate 5% SG, 8 gm per liter of water for control of American Army worm. Care should be taken that the maize crop which has been sprayed by insecticide should not be used as fodder.
<b>Chickpea</b>	Vegetative Growth Stage	During the flowering stage, irrigation should be applied to the crop as per availability. Since gram crop is very sensitive to water, if water is given more than required, the crop will be affected. To ensure proper availability of fertilizers to the crop and to increase the yield significantly, sprinkler irrigation method should be used.
<b>Banana</b>	Vegetative stage	As the minimum temperature is likely to be low, irrigation should be applied to the orchard in night hours. In the early morning make smoke in the orchard by burning green leaves. Apply 250 to 1000 grams of neem cake per tree to the banana plants depending on the crop stage. Cover the banana bunches with 6% perforated white plastic bag. Due to the present weather, for control of sigatoka disease, the affected leaf part/leaves should be removed and burned outside the orchard and the banana suckers should be cut by chopper periodically. Also, spray metiram 55% + pyraclostrobin 5% (60% WG) 30 grams per 10 liters of water. After that, 2 sprays should be done at an interval of 15 to 20 days.
<b>Animal Management</b>		Keep cattles inside the sheds during night and provide dry bedding to protect them from cold. Increase protein level and minerals in the feed concentrate to keep the animals healthy to cope up with the cold condition. Give animals mineral mixture along with salt regularly and also wheat grains, jaggery etc. 10%-20% in the daily ration during winter season to meet the energy requirement of the animals. Do not allow cattle/goats grazing during morning hours. Do not keep cattle/goat in the open during night time. In Poultry, keep the chicks warm by providing artificial light in the poultry sheds.
<b>Note: Farmers should take necessary precautionary measures while spraying insecticide, fungicide etc. and use Kisan Kavach Body Suite.</b>		

**Source:**

- 1) Weather Forecast : Regional Meteorological Centre (RMC, Mumbai)  
2) Last week weather summary : -

**Place : COA, Pune**

**Date : 05.12.2025**

**Sd/-**

**Principal Nodal Officer, GKMS, AMFU Pune &  
Head, Department of Agril. Meteorology, COA, Pune.**