

AGROMET ADVISORY BULLETIN GRAMIN KRISHI MAUSAM SEWA, AMFU, PUNE Department of Agricultural Meteorology College of Agriculture, Pune 411 005



Ph No. 020-25538009

E-mail: amfupune@gmail.com

Weather based Agromet Advisory committee meeting dated 08.10.2024 District Jalgaon

Weather Parameters			Weather Forecast (08.10.2024 to 12.10.2024)					
Date		9	10	11	12	13		
Rainfall (mm)		2	3	4	7	3		
Max. Temp. (0C)		36	35	34	34	35		
Min. Temp. (0C)		22	22	23	23	23		
Cloud Cover		1	6	6	8	4		
Max. RH (%)		73	75	72	74	75		
Min. RH (%)		44	43	43	46	48		
Wind Speed(km/hr)		8	7	8	6	10		
Wind direction (deg)		71	77	109	108	108		
Agromet Advisory Based			l on Weat	ther Fore	ecast Pre	diction		
Crop	Crop Stage	Advisory						
	mmary/ Alert	According to for						
		India Meteorological Department there is possibility of rainfall at isolated						
		places during dt. 08 th to 12 th October, 2024 in the district.						
	ange Forecast	As per ERFS products during 13 th to 19 th October, 2024 over Madhya						
(ERFS)		Maharashtra (Dhule, Nandurbar, Jalgaon, Nashik, Ahmednagar, Pune, Satara,						
		Sangli, Solapur, Kolhapur) division						
		Rainfall may remain above normal.						
		 Maximum temperature may remain normal. Minimum temperature may remain normal. 						
Concept A designment		Minimum temperature may remain normal.						
General Advisory		• As there is possibility of rainfall in the district spraying of insecticide and						
		pesticide should be done by considering the forecasted rain and preferably when there is no rain and use sticker while spraying						
		 when there is no rain and use sticker while spraying. Store the harvested produce at safe storage places. The harvested produce 						
		• Store the harvested produce at safe storage places. The harvested produce should be covered properly.						
		 Farmers should observe the pest and disease incidence on their field and if 						
		the infestation is above economic threshold level (ETL) then proper pest						
		and disease management should be done.						
		• Farmers should download and make use of Meghdoot mobile app for						
		weather based crop advisory and weather forecast.						
		• Also, farmers should download and make use of use of Damini mobile						
		app for lightni	ing and thund	derstorm war	ming and for	ecasting.		
		• Preparatory tillage operations should be done for rabi season crops for						
	1	sowing.						
Cotton	Picking stage	The picking of co						
		out. The cotton p		•				
		1.5 lakh eggs per	hectare of the	richogramma	a. For chemi	cal control	ot pink boll	

		worm spray profenophos 50 EC 30 ml or spinosad 45 SC 3.5 ml per 10 lit. of			
		water when there is no rain.			
Soybean		Harvesting of matured crop should be done when the colour of the pods			
	Maturity to	become yellowish red. If harvesting is delayed there is possibility of			
	harvesting	shattering of pods. The harvested produce should be kept in small heaps in			
	stage	the field itself to dry it in sunlight. After that carryout threshing operation. As there is possibility of rainfall at isolated places in the district the harvested			
		produce should be kept at safe storage places.			
Pigeon Pea		Apply irrigation to crop when it is in flowering stage as per availability and			
0		requirement. For control of various types of insect pests install Pheromone			
	Flowering	traps @ 5 traps per hectare. Hoeing or weeding should be done as required so			
	stage	that the competition between the crop and weeds for soil moisture and soil			
	~6	nutrients can be avoided. Erect "T" shape bird perches 50 to 60 per hectare in the field. As a propulsion array 5% near orby when the grap is in			
		the field. As a precautionary measure spray 5% neem ark when the crop is in the flowering.			
Maize	Maturity to	Care should be taken that the maize crop which have been sprayed by			
	harvesting	insecticide should not be feed as fodder to cattles at least for 15 to 20 days			
	stage	after spraying. Harvesting of the matured crop should be done and keep the			
		harvested produce in sunlight to dry. Store the harvested produce at safe			
		storage places. As there is possibility of rainfall in the district the harvested			
		produce should be covered properly.			
Chick pea	Sowing	Field should be prepared for the sowing of crop. Sowing of irrigated chick pea should be done between 20 th October to 10 th November for getting good			
		yield. Sowing should be done by giving seed treatment at proper depth and at			
		optimum soil moisture condition.			
Grapes	Pruning stage	October prunning should be carried out and the prunned branches and leaves			
•		should be destroyed properly. Apply paste of 0.4 % copper oxychloride + kav			
	<u> </u> '	(red soil) 3 kg per 10 lit. of water to the cut portion.			
Animal		For control and prevention of diseases all animals should be given medicine			
Managemen		and vaccination in consultation with veterinary doctor. During rainy season			
L		do not give excess of green fodder to animals. To prevent nuisance from ectoparasites spray Deltamethrin (Ex. Butox) by taking advice from			
		Veterinary doctor in goat shade and on the goats. Drinking water tanks should			
		be cleaned regularly and should be applied lime.			
Note: Farmers should take necessary precautionary measures while spraying insecticide, fungicide etc.					

Source:

1) Weather Forecast	: Regional Meteorological Centre (RMC, Mumbai)
---------------------	--

2) Last week weather summary : -

Place : COA, Pune Date : 08.10.2024 Sd/-Principal Nodal Officer, GKMS, AMFU Pune & Head, Department of Agril. Meteorology, COA, Pune.