



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, PUNE
Department of Agricultural Meteorology
College of Agriculture, Pune 411 005



Ph No. 020-25538009

E-mail: amfupune@gmail.com

Weather based Agromet Advisory committee meeting dated 23.09.2022
District Jalgaon

Weather Parameters	Weather Forecast (23.09.2022 to 27.09.2022)				
Date	24	25	26	27	28
Rainfall (mm)	14	6	4	14	15
Max. Temp. (0C)	28	31	32	32	33
Min. Temp. (0C)	22	23	24	24	25
Cloud Cover	7	5	4	6	5
Max. RH (%)	89	90	89	89	88
Min. RH (%)	64	63	56	59	59
Wind Speed(km/hr)	17	14	13	15	14
Wind direction (deg)	270	270	283	291	294

Agromet Advisory Based on Weather Forecast Prediction

Crop	Crop Stage	Advisory
Weather Summary/ Alert		According to forecast given by Regional Meteorological Centre, Mumbai of India Meteorological Department there is possibility of rainfall at some (scattered) places on dt. 23 rd & 24 th September, 2022 and at isolated places on dt. 25 th , 26 th & 27 th September, 2022 in the district.
Extended Range Forecast (ERFS)		As per ERFS products during 25 th September, 2022 to 01 st October, 2022 over Madhya Maharashtra (Dhule, Nandurbar, Jalgaon, Nashik, Ahmednagar, Pune, Satara, Sangli, Solapur, Kolhapur) division <ul style="list-style-type: none"> ➤ Rainfall may remain above normal. ➤ Maximum temperature may remain normal. ➤ Minimum temperature may remain normal.
General Advisory		<ul style="list-style-type: none"> • Spraying of insecticide and pesticide should be done by considering the forecasted rain and preferably when there is no rain. • Farmers should download and make use of Meghdoot mobile app for weather based crop advisory and weather forecast. • Also, farmers should download and make use of use of Damini mobile app for lightning and thunderstorm warning and forecasting. • Farmers should observe the pest and disease incidence on their field and if the infestation is above economic threshold level (ETL) then proper pest and disease management should be done. • The harvested kharif crops should be kept at safe storage places.
Cotton	Flower bud to flowering stage	Water logging condition in cotton crop may lead to incidence of wilt. For its control drench the mixture of 1.5 kg Urea + 1.5 kg Potash in 100 liter of water and pour 150 to 200 ml per plant. After 8-10 days mix 2 kg DAP in 100 liters of water and pour 150 to 200 ml per plant. Due to present cloudy and humid condition, cotton crop may be protected from sap sucking pests by spraying 5% neem ark when there is no rain. The cotton plant may get

		infested by green leaf eating caterpillar, for its control erect 5 pheromone traps per hectare. The cotton plant infested by Pink bollworm, for its control disseminate 1.5 lakh eggs per hectare of trichogramma. For chemical control of pink boll worm spray profenophos 50 EC 30 ml or spinosad 45 SC 3.5 ml per 10 lit. of water when there is no rain.
Sugarcane	Vegetative growth stage	For control of white grub infestation apply 20 kg of metarhizium anisoplie per hectare in soil or 2.5 to 3 kg per hectare for spraying.
Soybean	Pod development	Erect 8 to 10 bird perches per acre for control of various types of larval pest. For control of leaf eating spodoptera larva spray erect 5 spodulure pheromone traps per hectare, collect and destroy the egg masses and larva which will help in controlling the pest. If chemical control is required then spray 20% Chloropyriphos 20 ml or Cypermethrine 25 %, 4 ml per 10 liter of water. High humidity and cloudy condition is favourable for occurrence of rust in soybean. For control of rust disease spray propiconyazole or hexaconyazole 10 ml per 10 litre of water.
Pigeon Pea	Branching stage	For control of various types of insect pests install Pheromone traps @ 5 traps per hectare. Care should be taken to avoid water logging condition in standing crop by draining out excess rainwater. Hoeing or weeding should be done as required so that the competition between the crop and weeds for soil moisture and soil nutrients can be avoided.
Green gram & black gram	Pod development to maturity stage	Start harvesting immediately when 75 % of pods mature and when there are no rains. One or two rounds of pod picking should be done after 8-10 days to avoid shattering of pods and avoid losses. The harvested crop should be kept at safe storage places. After proper drying threshing should be carried out. The threshed grains should be dried for 5-6 days before filling in the bags. Storage of these bags should not be done in moist place. Mix neem leaves (5%) in the grains to avoid storage infestation.
Maize	Flowering stage	If the infestation of Fall Army Worm is more than 10% for its control take spray of Chlorantriliprole 18.5 SC 4 ml per 10 lit. of water when there are no rains. For prevention of infestation of cob borer spread Trichogramma chilonis egg parasitoids in the field at silking stage.
Brinjal	Vegetative growth stage	Due to present cloudy and humid weather condition the infestation of fruit and shoot borer may occur for its control chlorantriliprol 18.5% S.C., 4 ml or chlorpyriphos 20% EC, 20 ml per 10 liters of water in a clean and clear weather condition.
Mango		Due to present cloudy and humid condition for control of various insect pest and diseases on new shoots of mango plants spray 5% lymda psyhalothrin 6 ml or quinolphos 25 ml + 20gram carbendenzim per 10 lit. of water when there is no rain.
Banana		Due to present cloudy and humid weather the incidence of blast (sigatoka) disease may occur and for its control destroy the part of leaf or full leaf of plants. When the incidence is noticed spray propiconyazol 5 ml and mineral oil 100 ml per 10 liter of water when there are no rains. Repeat two spray at an interval of 20 days.
Animal Management		To avoid the spreading of diseases in cattle's care should be taken to isolate the infected animal and the excreta of the animal should be destroyed by burying it far away. Goat pox vaccine should be administered on the advice of a veterinarian to control lumpy skin disease. Due to open grazing the animals feed on grasses grown in the monsoon and there digestive system gets disturbed and they may suffer by Diarrhoea. So the proportion of dry fodder feed should be increased comparatively during rainy season. It is most

		important to give wormicide medicine to animals by consulting veterinary doctor. Always give clean drinking water to animals. For control and prevention of diseases all animals should be given vaccine and medicine in consultation with veterinary doctor.
--	--	---

Note: Farmers should take necessary precautionary measures while spraying insecticide, fungicide etc.

Source:

- 1) Weather Forecast : Regional Meteorological Centre (RMC, Mumbai)
 2) Last week weather summary : -

Place : COA, Pune

Date : 23.09.2022

Sd/-

**Principal Nodal Officer, GKMS, AMFU Pune &
 Head, Department of Agril. Meteorology, COA, Pune.**