



# AGROMET ADVISORY BULLETIN

## GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR

### ZONAL AGRICULTURAL RESEARCH STATION, SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: [adrkolhapur@rediffmail.com](mailto:adrkolhapur@rediffmail.com)

## Weather based Agromet Advisory committee meeting dated 30.05.2025

District: SANGLI

Last Week Weather Summary (24.05.2025 to 30.05.2025)							Weather Parameters	Weather Forecast (31.05.2025 to 04.06.2025)				
24	25	26	27	28	29	30	Date	31	01	02	03	04
-	-	-	-	-	-	-	Rainfall (mm)	3	3	5	7	3
-	-	-	-	-	-	-	Max. Temp. (°C)	30	31	32	32	33
-	-	-	-	-	-	-	Min. Temp. (°C)	24	24	24	25	25
-	-	-	-	-	-	-	Cloud Cover	5	6	7	7	5
-	-	-	-	-	-	-	Max. RH (%)	83	81	77	80	81
-	-	-	-	-	-	-	Min. RH (%)	82	73	71	76	72
-	-	-	-	-	-	-	Wind Speed(km/hr)	24	25	23	23	22
-	-	-	-	-	-	-	Wind Direction	W	WSW	WSW	WSW	WSW

### Agromet Advisory Based on Weather Forecast

Crop	Stage	Advisory
Weather Summary		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is forecast of generally cloudy sky from 31 <sup>st</sup> May to 4 <sup>th</sup> June in Sangli district. <b>There is forecast of light rainfall at isolated places from 31<sup>st</sup> May to 4<sup>th</sup> June. During next 5 days Tmax may remain between 30.0 to 33.0°C. During next 5 days the minimum temperature may remain nearby 24.0 to 25.0°C.</b> Morning relative humidity will remain nearby 77 to 83% and Afternoon relative humidity will remain nearby 71 to 82%. Wind speed will remain between 22 to 25 km per hour.
ERFS		According to Extended Range Forecast (ERF) in Madhya Maharashtra Sub Division rainfall will remain below normal, maximum temperature will remain Moderately Below Normal and minimum temperature will remain below normal valid from 4 <sup>th</sup> to 10 <sup>th</sup> June, 2025
General Advisory		➤ Carry out the preparatory tillage operation like ploughing for cultivation of crop in Kharif season. For taking pulse crop in Kharif season carry out deep ploughing followed by 2 harrowing, mix Compost @ 5 ton per hector before last harrowing which will help in better mixing of compost in soil.
SMS Advisory		Purchase Improved/Hybrid seed from government authorised centres or from Sell units of Agriculture University.
Maize	Presowing	➤ Land : Select Medium to heavy, well drained especially riverside alluvial soil, having well organic matter content and having well water holding capacity. ➤ Land preparation: Carry out one deep ploughing (15 to 20 cm) and 2 to 3 harrowing. Mix well decomposed FYM or Compost @ 10 to 12 ton per hector at the time of last harrowing.
Soybean	Land preparation	➤ Land: Select Medium and well drained soil ➤ Land preparation: One ploughing followed by 2 harrowing. ➤ Mix well decomposed FYM or Compost @ 10 to 12 ton per hector at the time of last harrowing. ➤ Make arrangements of quality and certified seeds for sowing of crops in the Kharif season.
Groundnut	Land preparation	➤ Land: Select Medium and well drained soil ➤ Land preparation: One ploughing followed by 2 harrowing. ➤ Mix well decomposed FYM or Compost @ 10 ton per hector before last harrowing. ➤ Make arrangements of quality and certified seeds for sowing of crops in the Kharif season.
Tomato	Sowing/Nursery	➤ For Kharif season Tomato cultivation carry out sowing of Tomato seeds on raised beds of Size 3m length, 1m width and 15 cm height. Removes stones and soil clods from beds and mix 1 ghamela of well decomposed FYM, 50 gram Copper Sulphate and 100 to 150 gram Suphala and make bed plain. Make parallel lines on beds with 7 to 8cm distance and carry out sowing of seeds in that lines. Cover these lines with soil and give water to bed with watering can. The plants will ready to transplant within 3 to 5 weeks. ➤ Use Bhagyashree, Dhanashree varieties for sowing. For Long distance market: Phule Raja (Hybrid), More Beta Carotene: Phule Kesari and Cherry Tomato: Phule Jayshree. ➤ Carry out nursery preparation only if there is assured source of irrigation.



**AGROMET ADVISORY BULLETIN**  
**GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR**  
**ZONAL AGRICULTURAL RESEARCH STATION,**  
**SHENDA PARK, KOLHAPUR**



Ph. No. 0231 2692416

E-mail: [adrkolhapur@rediffmail.com](mailto:adrkolhapur@rediffmail.com)

<b>Sugarcane</b>	<b>Establishment</b>	<p>➤ <b>White grub</b> : Adults comes together on Acacia, Neem and Ber trees at the time of evening after first shower of rain. First female adults and then male adults come out of the soil. They eat leaves of these trees. During night period shake the branches of these trees due to which adults falls on ground. Collect these adults and destroy them in kerosene mixed water.</p>
<b>Ginger</b>	<b>Planting</b>	<p>➤ Ginger planting should be done up to the first week of June.</p> <p>➤ For planting, select improved varieties like Varada, Rijatha, Mahim, and Mahima.</p> <p>➤ Rhizomes should be planted 4 to 5 cm deep. Ensure that the rhizomes are completely covered with soil. Ginger requires moderate sunlight.</p> <p>➤ According to the slope of the land, decide the method of planting and prepare beds in the direction of the slope. Make sure the beds are uniformly level.</p> <p>➤ While preparing beds, use the tractor's furrow opener to make straight beds. During planting, take out the seed rhizomes from storage and cut them into uniform pieces (known as "modes").</p> <p>➤ The day the ginger piece is planted in the soil, it must receive ample water. Water should be given so that moisture penetrates at least 4 to 5 inches below the planted piece. If this is not achieved, water again the next day to ensure proper soil moisture. Subsequent irrigation should be given every alternate day or every two days depending on soil type, maintaining constant soil moisture.</p> <p>➤ About eight days after planting, give a light covering of soil (earthing up), as the tips of the longer pieces planted are usually at soil surface level. This is necessary to protect them from heat and prevent exposed buds from getting damaged.</p> <p>➤ After planting, it is important to control the soil temperature by irrigation. Soil temperature is generally higher compared to the surrounding air. Another important point is to maintain the moisture content in the seed rhizomes. If the rhizomes lose moisture, it affects germination and the growth of new shoots.</p>
<b>Turmeric</b>	<b>Cultivation</b>	<p>➤ If there is possibility of irrigation carry out cultivation of turmeric upto first week of June</p> <p>➤ Use Phule Swarupa, Selem, Krushna, Rajapuri and Tekurpeta varieties for cultivation.</p> <p>➤ Use 25 to 30 quintals of rhizome seed per hectare cultivation.</p> <p>➤ Intercrop- Turmeric+Frenchbean, Turmeric+ Fenugreek, Turmeric+ Chilli</p> <p>➤ Fertilizer- Apply 25 to 40 ton of well decomposed Organic Manure and 100kg Phosphorus and 100 kg Potash per hectare.</p> <p>➤ Organic Turmeric</p> <p>➤ Biological Rhizome Treatment: Prepare mixture of 500 liters of water+ VAM 12.5 Kg +PSB 5 kg+ Azospirillum 5 kg and Dip rhizomes in it for 15 minutes.</p> <p>➤ Organic Manures: FYM 15-20 ton/ha + Neem Seed Kernel Extract 4 ton/ha + Vermicompost 2 ton/ha.</p>
<b>Animals</b>	<b>Growth</b>	<p>➤ Carry out vaccination to animals by taking consultation of Veterinarian for prevention of diseases like Foot and Mouth Disease, Black Quarter and Hemorrhagic Septicemia.</p>
<b>Poultry</b>	<b>Growth</b>	<p><b>Care of Poultry Birds During the Rainy Season</b></p> <ol style="list-style-type: none"><li><b>Shelter and Housing:</b><ul style="list-style-type: none"><li>○ Ensure the poultry shed is <b>well-roofed and leak-proof</b>.</li><li>○ Maintain <b>dry and clean flooring</b> by using adequate bedding (like dry straw or sawdust).</li><li>○ Elevate the shed floor slightly or improve drainage to <b>prevent waterlogging</b>.</li><li>○ Protect against <b>cold drafts and dampness</b>, which can stress the birds.</li></ul></li><li><b>Ventilation and Hygiene:</b><ul style="list-style-type: none"><li>○ Maintain <b>proper ventilation</b> without exposing birds to direct cold winds.</li><li>○ Clean and disinfect the coop regularly to <b>prevent fungal and bacterial infections</b>.</li><li>○ Remove wet litter and replace it frequently to avoid <b>respiratory issues</b> and <b>foot infections</b>.</li></ul></li><li><b>Feed and Water:</b><ul style="list-style-type: none"><li>○ Store feed in a <b>dry, moisture-proof container</b> to prevent fungal growth</li><li>○ Provide <b>clean, fresh drinking water</b>; check for contamination after heavy rains.</li><li>○ Add <b>vitamins and electrolytes</b> to water, especially Vitamin C and A, to help boost immunity.</li></ul></li><li><b>Health and Disease Management:</b><ul style="list-style-type: none"><li>○ Watch for signs of diseases like <b>coccidiosis, fowl cholera</b>, and respiratory infections.</li><li>○ Ensure regular <b>vaccination and deworming</b> as per veterinary guidance.</li><li>○ Isolate any sick birds immediately to avoid spread.</li></ul></li></ol>





**AGROMET ADVISORY BULLETIN**  
**GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR**  
**ZONAL AGRICULTURAL RESEARCH STATION,**  
**SHENDA PARK, KOLHAPUR**



Ph. No. 0231 2692416

E-mail: [adrkolhapur@rediffmail.com](mailto:adrkolhapur@rediffmail.com)

	<p>5. <b>Mosquito and Parasite Control:</b></p> <ul style="list-style-type: none"><li>○ Clear stagnant water around the shed to <b>prevent mosquito breeding</b>.</li><li>○ Use safe insect repellents or mosquito nets if necessary.</li><li>○ Dust birds and bedding with appropriate <b>anti-parasitic powders</b> to control lice and mites.</li></ul>
<p>To download <b>“Meghdoot”</b> app scan QR code</p> 	<p>To download <b>“Damini”</b> app scan QR code</p> 

**Source:**

1) Weather Forecast

: Research Section, Mumbai

**Place : ZARS, KOLHAPUR**

**Date : 30.05.2025**

**Sd/-**

**Nodal Officer, GKMS, &  
Associate Director of Research,  
ZARS, Kolhapur**