



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

Weather based Agromet Advisory committee meeting dated 05.12.2025

District: SATARA

Last Week Weather Summary (29.11.2025 to 05.12.2025)							Weather Parameters	Weather Forecast (06.12.2025 to 10.12.2025)				
29	30	01	02	03	04	05	Date	06	07	08	09	10
-	-	-	-	-	-	-	Rainfall (mm)	0	0	0	0	0
-	-	-	-	-	-	-	Max. Temp. (°C)	31	30	30	29	29
-	-	-	-	-	-	-	Min. Temp. (°C)	17	16	16	15	14
-	-	-	-	-	-	-	Cloud Cover	5	4	2	0	0
-	-	-	-	-	-	-	Max. RH (%)	75	68	64	59	64
-	-	-	-	-	-	-	Min. RH (%)	58	51	46	41	42
-	-	-	-	-	-	-	Wind Speed(km/hr)	10	9	7	6	8
-	-	-	-	-	-	-	Wind Direction	ENE	ENE	ENE	ENE	ENE

Agromet Advisory Based on Weather Forecast Prediction

Crop	Stage	Advisory
Weather Summary/ Alert		Weather forecast given by Regional Meteorological Centre, Mumbai has indicated that there is forecast of generally cloudy sky on 6 th and 7 th December and forecast of mainly clear sky on 8 th , 9 th and 10 th December in Satara district. During next 5 days Tmax may remain 29.0 to 31.0°C and Tmin may remain between 14.0 to 17.0°C. Morning relative humidity will remain nearby 59 to 75% and Afternoon relative humidity will remain nearby 42 to 58%. Wind speed will remain between 6 to 10 km per hour.
ERFS		According to Extended Range Forecast (ERF) in Madhya Maharashtra Sub Division there will be no rainfall, maximum temperature will remain moderately below normal and minimum temperature will remain below normal valid from 10 th to 16 th December, 2025
General Advisory		<p>Due to the increasing cold, citizens should take the following precautions:</p> <ul style="list-style-type: none"> ➤ Take special care of elderly people and young children. ➤ Stay indoors as much as possible and avoid unnecessary travel. ➤ Wear sweaters, mufflers, ear caps, socks, and gloves. ➤ Regularly consume warm beverages. ➤ Stay updated on weather conditions by listening to the radio, watching TV, or reading newspapers. <p>As the intensity of cold increased farmers should take care of their crop in following way</p> <ul style="list-style-type: none"> ➤ To protect the crop from cold, provide light irrigation in the evening. ➤ Cover banana bunches with perforated polyethylene bags. ➤ Harvest workers should take special care. Avoid working early in the morning during cold warnings, if possible
SMS Advisory		➤ There is possibility of drop in minimum temperature by 2 to 3°C in upto 10 th December thus farmers should take care of themselves and their cattles
Rabbi Sorghum	Vegetative	<ul style="list-style-type: none"> ➤ The temperature dropped below 15°C thus the incidence of aphids is observed in some parts. ➤ Nymphs and Adults of Aphids sucks the sap from leaves and excretes honey like sugary substance on leaves thus black mould develops on this substance (Sticky/Sugary disease) which alters the food formation of plants and results in decrease in yield of crop. ➤ For control use yellow sticks traps @ 4 per hector and if infestation notices spray the crop with 5% Neem Seed Kernal Extract @ 5ml per liter of water.
Wheat	Beginning of tillering	<ul style="list-style-type: none"> ➤ If an infestation of aphids is observed on wheat crops (10 nymphs or adult aphids per branch per plant), for their control using biological components, apply Metarhizium anisopliae 1.15 WP at 40 grams or Verticillium lecanii 1.15 WP at 40 grams per 10 liters of water. Two sprays should be conducted at an interval of 10 to 15 days. ➤ Due to the cloudy weather there is possibility of incidence of Rust on Wheat plant. To control rust spray the crop with Mancozeb @ 1.5 kg by mixing in 500 liters of water per hector.



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

Chick Pea	Vegetative	➤ Chick pea crop is very sensitive to the excess irrigation, as excess irrigation causes wilting of plants. With the use of Sprinkler irrigation we can give irrigation as per the requirement of the crop and at any time which also maintain <i>vafasa</i> condition in soil which further leads to increase in crop yield thus Give irrigation to Chickpea crop through Sprinkler irrigation.
Sugarcane	Growth	➤ December Rate of evaporation increases to 3.63 mm/day thus Sugarcane needs 2.17 liters water per day so to fulfill this water requirement Drip Irrigation Unit require to run for 33 minutes daily.
Strawberry	Growth	➤ Due to high relative humidity there is possibility of sucking pest thus to control pest infestation use Yellow and Blue sticky traps. ➤ After 30 days of planting, the beds should be weeded and covered with black or silver-colored plastic mulching paper
Grape	Shoot Growth stage	1. In case the soil is under wapsa (field capacity) condition, donot irrigate the vineyard. 2. During shoot growth stage (Fruit pruning season), apply irrigation through drip @ 5950 - 8500 L/ acre/ day for all grape growing regions. In case vigour is more than desired, then reduce irrigation water application by half to 3000-4000 L/acre and stop nitrogen application. Still if growth is more, stop the irrigation till such time the growth is brought under control and then start irrigation. 3. From flowering to fruit setting, apply irrigation through drip upto 2000-2500 L/ acre/ day. Vigour needs to be controlled. 4. Practice mulching to keep the bunds moistened. This will reduce the salinity build up in the root zone due to evaporation of the moisture from the surface of the bund. 5. During Berry development stage, apply irrigation through drip @ 5950 - 8500 L/ acre/ day for all grape growing regions.
Brinjal	Fruit development	➤ Due to fluctuations in maximum and minimum temperatures during the past week, as well as high humidity in the air, an infestation of fruit and shoot borer has been observed in brinjal. Initially, the larva bores into the tender shoots when the plant has no fruits and feeds on the inner tissues, causing the shoots to dry up. Later, after fruit formation, the larvae bore into the fruits and feed inside, rendering the fruits unfit for market sale. ➤ The pest can cause 40–50% damage to the fruits. ➤ For control: About 20 days after planting, if damaged shoots and fruits are noticed, they should be collected and destroyed or buried in a deep pit. Spray 5% neem seed extract or Spinetoram 11.70 SC @ 5–7 ml, or Chlorpyrifos 20% EC @ 20 ml, or Spinosad 45% SC @ 4 ml per 10 litres of water using a hand sprayer. ➤ Install 40 pheromone traps (funnel type) per acre in the field for monitoring and management of the shoot and fruit borer.
Okra	Fruit development	➤ Due to cloudy and humid weather in last week there is incidence of Powdery mildew on Okara thus to control spray the crop with Carbendezium 10 gram or Hexaconazole 10 ml per 10 liters of water.
Turmeric	Rhizome Development	➤ Due to humid weather in last week there is possibility of leaf spot disease on Turmeric plant. Take the spray of Carbendezium 50W.P.10 gram by mixing in 10 liters of water. Use sticker while spraying.
Onion	Bulb Development	➤ Due to cloudy and humid weather in last week the incidence of Blight is observed on Onion thus to control spray the crop with Diphenconazole 25% E.C. 10 ml per 10 liters of water.
Animals	Growth	➤ Keep cattle's inside the sheds during night and provide dry bedding to protect them from cold. ➤ Increase protein level and minerals in the feed concentrate to keep the animals healthy to cope up with the cold condition. ➤ Do not keep cattle/goat in the open during night time.
Poultry	Growth	➤ There is forecast of Tmin to remain 11.0 to 14.0°C thus to protect poultry bird from clod keep the poultry shed warm particularly during night and early in morning.



AGROMET ADVISORY BULLETIN
GRAMIN KRISHI MAUSAM SEWA, AMFU, KOLHAPUR
ZONAL AGRICULTURAL RESEARCH STATION,
SHENDA PARK, KOLHAPUR



Ph. No. 0231 2692416

E-mail: adrkolhapur@rediffmail.com

To download “**Meghdoot**” app scan QR code



To download “**Damini**” app scan QR code



Source:

1) Weather Forecast : Research Section, Mumbai

Place : ZARS, KOLHAPUR

Date : 05.12.2025

Sd/-

Nodal Officer, GKMS, &
Associate Director of Research,
ZARS, Kolhapur